

SURAT KETERANGAN

Nomor: 181/UNUSA/Adm-LPPM/III/2019

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 15 Maret 2019.

Judul : Predictors For The Treatment Effect Of Sodium Glucose Co-Transporter 2 Inhibitors In Patients With Type 2 Diabetes Mellitus

Penulis : Shusuke Yagi, Ken-ichi Aihara, Takeshi Kondo, Kiyoe Kurahashi, Sumiko Yoshida, Itsuro Endo, Daiju Fukuda, Yutaka Nakaya, Kin-ichiro Suwaki, Takashi Takeji, Toshihiro Wada, Hotimah Masdan Salim, Saori Hama, Tomomi Matsuura, Takayuki Ise, Kenya Kusunose, Koji Yamaguchi, Takeshi Tobiume, Hirotsugu Yamada, Takeshi Soeki, Tetsuzo Wakatsuki, Munehide Matsuhisa, Michio Shimabukuro, Masashi Akaike, Masataka Sata

Identitas : Advances in therapy, Vol. 35, No.1, 124-134, Tahun 2018

No. Pemeriksaan : 2019.03.15.68

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 24%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya

Surabaya, 15 Maret 2019

Ketua LPPM,



Dr. Istas Pratomo, S.T., M.T.

NPP. 16081074

paper 4

by Hotimah Dr

Submission date: 15-Mar-2019 02:52PM (UTC+0700)

Submission ID: 1093757441

File name: Predictors_1.pdf (817.79K)

Word count: 4990

Character count: 27290

ORIGINAL RESEARCH

Predictors for the Treatment Effect of Sodium Glucose Co-transporter 2 Inhibitors in Patients with Type 2 Diabetes Mellitus

Shusuke Yagi · Ken-ichi Aihara · Takeshi Kondo · Kiyoe Kurahashi · Sumiko Yoshida · Itsuro Endo · Daiju Fukuda · Yutaka Nakaya · Ken-ichi Suwaki · Takashi Takeji · Toshihiro Wada · Hotimah Masdan Salim · Saori Hama · Tomomi Matsuura · Takayuki Ise · Kenya Kusunose · Koji Yamaguchi · Takeshi Tobiume · Hirotugu Yamada · Takeshi Soeki · Tetsuzo Wakatsuki · Munehide Matsuhisa · Michio Shimabukuro · Masashi Akaike · Masataka Sata

Received: August 6, 2017

© Springer Healthcare Ltd., part of Springer Nature 2017

ABSTRACT

Introduction: Predictors for the effect of sodium glucose co-transporter 2 (SGLT2) inhibitors at lowering hemoglobin A1c (HbA1c) levels in type 2 diabetes mellitus patients remain unclear. We therefore aimed to elucidate the predictors in type 2 diabetes patients after 3 months of SGLT2 treatment.

Methods: A total of 302 consecutive type 2 diabetes patients had been treated with SGLT2 inhibitors as monotherapy or add-on therapy to existing antidiabetic treatments were

enrolled retrospectively. After excluding 27 patients whose HbA1c levels could not be evaluated 3 months after treatment, the glucose-lowering effects of SGLT2 inhibitors were assessed in 275 patients by measuring HbA1c levels before and 3 months after treatment. The predictors for changes in HbA1c levels after 3 months of treatment were evaluated.

Results: SGLT2 inhibitor treatment for 3 months decreased HbA1c levels from $7.8 \pm 1.2\%$ to $7.4 \pm 1.0\%$ ($p < 0.0001$). A multiple regression analysis showed that the independent determinants for SGLT2 inhibitor treatment effect included decreased HbA1c levels after 1 month of treatment, high baseline HbA1c levels, and a high estimated glomerular filtration rate (eGFR).

Conclusion: We show that type 2 diabetes patients who received the greatest glucose-lowering effect with SGLT2 inhibitor treatment were those with preserved renal function (high

Enhanced content To view enhanced content for this article go to <http://www.medengine.com/Redeem/7DCCF0600B76BF39>.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s12325-017-0639-z>) contains supplementary material, which is available to authorized users.

S. Yagi (✉) · H. M. Salim · S. Hama · T. Matsuura · T. Ise · K. Kusunose · K. Yamaguchi · T. Tobiume · Hirotugu Yamada · T. Soeki · T. Wakatsuki · M. Sata
Department of Cardiovascular Medicine,
Tokushima University Graduate School of
Biomedical Sciences, Tokushima, Japan
e-mail: syagi@tokushima-u.ac.jp

S. Yagi
Department of Community Medicine and Human
Resource Development, Tokushima University
Graduate School of Biomedical Sciences,
Tokushima, Japan

S. Yagi · Y. Nakaya · K. Suwaki · T. Takeji · T. Wada
Department of Internal Medicine, Shikoku Central
Hospital, Shikokuchuo, Ehime, Japan

K. Aihara
Department of Community Medicine for Diabetes
and Metabolic Disorders, Tokushima University
Graduate School of Biomedical Sciences,
Tokushima, Japan

baseline eGFR) and high baseline HbA1c levels. Moreover, SGLT2 inhibitor treatment efficacy could be predicted by the patients' initial response to treatment.

Keywords: Estimated glomerular filtration rate; Glucose-lowering; HbA1c; Predictors; Type 2 diabetes mellitus

INTRODUCTION

Selective sodium glucose co-transporter 2 (SGLT2) inhibitors improve glucose metabolism by inhibiting SGLT2, which is expressed in the proximal tubule of the kidney. Thus, SGLT2 inhibitors enhance the renal excretion of glucose and decrease blood glucose levels in patients with type 2 diabetes [1].

The EMPA-REG OUTCOME trial has shown a significant reduction in cardiovascular-related and all-cause mortality in patients with type 2 diabetes, who are at high risk for cardiovascular events, when the SGLT2 inhibitor empagliflozin was added to their standard care [2]. The pleiotropic effects of SGLT2 inhibitors, including lower blood pressure, body weight, visceral fat content, uric acid, albuminuria, increased

high-density lipoprotein (HDL) cholesterol, and improved endothelial function might be involved in the improvement of cardiovascular outcomes [3, 4].

There are some patients whose HbA1c levels do not respond to SGLT2 inhibitors therapy, or whose HbA1c levels rebound after an initial decrease in response to SGLT2 inhibitors. If we can predict patients who show a significant response to treatment with SGLT2 inhibitors, it would be of great clinical value; however, few studies have distinguished the patients who show the greatest glucose-lowering effect upon treatment with SGLT2 inhibitors.

In this study, we aimed to elucidate the predictors for changes in HbA1c levels after 3 months of treatment in a clinical setting.

METHODS

Study Population

SGLT2 inhibitors (ipragliflozin, dapagliflozin, luseogliflozin, tofogliflozin, canagliflozin, or empagliflozin) were administered to 302 consecutive type 2 diabetic patients as monotherapy or as an add-on therapy to existing antidiabetic treatments at the Department of Cardiovascular Medicine and Department of Endocrinology and Metabolism at the Tokushima University Hospital, Japan, and the Department of Internal Medicine at the Shikoku Central Hospital, Japan, from July 2014 to August 2016.

A total of 27 patients were excluded from the analysis as we could not evaluate their HbA1c levels 3 months after treatment due to several reasons, including discontinuation of treatment or minor adverse effects of the SGLT2 treatment (Fig. 1). No patients experienced severe side effects such as hypoglycemic episodes or diabetic ketoacidosis resulting in a hospital visit or hospitalization during the 3-month SGLT2 inhibitor treatment period.

We enrolled 275 type 2 diabetes patients and evaluated the glucose-lowering effects of SGLT2 inhibitors (Fig. 1).

T. Kondo · K. Kurahashi · S. Yoshida · I. Endo
Department of Hematology, Endocrinology and Metabolism, Tokushima University Graduate School of Biomedical Sciences, Tokushima, Japan

D. Fukuda · M. Shimabukuro
Department of Cardio-Diabetes Medicine, Tokushima University Graduate School of Biomedical Sciences, Tokushima, Japan

M. Matsuhisa
Diabetes Therapeutics and Research Center, Institute of Advanced Medical Sciences, Tokushima University, Tokushima, Japan

M. Shimabukuro
Department of Diabetes, Endocrinology and Metabolism, School of Medicine, Fukushima Medical University, Fukushima, Japan

M. Akaike
Department of Medical Education, Tokushima University Graduate School of Biomedical Sciences, Tokushima, Japan




Fig. 1 Flow chart of the study

Measures

The glucose-lowering effects of the SGLT2 inhibitors (HbA1c at 3 months—baseline HbA1c) were retrospectively assessed in 275 patients by measuring HbA1c levels (National Glycohemoglobin Standardization Program reference values) before and 3 months after treatment. The proportion of patients who achieved glycemic control of the HbA1c target of <7%, recommended by the guideline to reduce complications of diabetes, was also calculated [5]. In addition to HbA1c levels, we assessed the patients' serum lipid profiles [including low-density lipoprotein (LDL) cholesterol, HDL cholesterol, and triglyceride levels]; renal function, including serum creatinine levels and estimated glomerular filtration rates (eGFR); hematocrit; body weight; and blood pressure. Predictors for changes in HbA1c levels after 3 months of treatment were evaluated by simple and multiple regression analysis.

All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1964, as revised in 1983. The study protocol was approved by the Ethics Committee of Tokushima University Hospital (No. 2432) and Shikoku Central Hospital (No. 25).

Statistical Analysis

We calculated the averages for continuous variables. Values are expressed as means \pm standard deviations and percentages for continuous and categorical parameters, respectively. Seasonal changes in HbA1c levels were evaluated by analysis of variance (ANOVA), and changes in HbA1c levels with/without concomitant antidiabetic drug treatment were evaluated using the unpaired *t* test and multiple regression analysis. HbA1c levels, blood glucose levels, and other clinical parameters before and 3 months after treatment with SGLT2 inhibitors were compared using the paired *t* test. The degrees of association among independent variables for HbA1c levels after 3 months of treatment were assessed by multiple regression analysis. All statistical analyses were performed using the JMP software (v.11; SAS Institute, Cary, NC, USA). Statistical significance was defined as $p < 0.05$.

RESULTS

Clinical Characteristics of the Study Participants

The characteristics of the patients enrolled in this study are shown in Table 1. They had moderate diabetes with a mean HbA1c level of $7.8 \pm 1.2\%$ and were obese with a mean body mass index (BMI) of $31 \pm 7 \text{ kg/cm}^2$. Their mean eGFR was $83 \pm 26 \text{ mL/min/1.73 m}^2$.

Effects of SGLT2 Inhibitor Treatment on HbA1c Levels

Three months of treatment with SGLT2 inhibitors led to a significant decrease in HbA1c levels (HbA1c at 3 months—baseline HbA1c: $-0.4 \pm 0.9\%$, $p < 0.0001$; Table 2).

In the analysis of seasonal changes in HbA1c levels 3 months after the start of SGLT2 inhibitors treatment, we found that HbA1c levels were higher in February and March when compared to other months; however, this was not statistically significant (Fig. 2).

Table 1 Clinical characteristics of study subjects

Variables	
Number of patients	275
Male/female	153 (56%)/122 (44%)
Age (years)	54 ± 13
Body mass index (kg/m ²)	31 ± 7
HbA1c (%)	7.8 ± 1.2
Casual blood glucose (mg/dL)	167 ± 57
LDL cholesterol (mg/dL)	113 ± 33
Triglyceride (mg/dL)	212 ± 297
HDL cholesterol (mg/dL)	51 ± 15
Hematocrit (%)	41 ± 5
Serum creatinine (mg/dL)	0.76 ± 0.39
eGFR (ml/min/1.73 m ²)	83 ± 26
Systolic blood pressure (mmHg)	134 ± 16
Diastolic blood pressure (mmHg)	80 ± 12
Co-morbidities	
Hypertension	162 (59%)
Dyslipidemia	141 (51%)
Coronary artery disease	28 (10%)
Stroke	3 (1%)
Antidiabetic drugs	
SGLT2 inhibitors, mean dose	
Ipragliflozin	38 (14%), 45 ± 10 mg
Dapagliflozin	71 (26%), 5.5 ± 1.5 mg
Luseogliflozin	50 (18%), 2.7 ± 0.6 mg
Tofogliflozin	51 (19%), 20 ± 0 mg
Canagliflozin	50 (18%), 100 ± 0 mg
Empagliflozin	15 (5%), 12 ± 5 mg
Other anti-diabetic drugs	
DPP-4 inhibitors	123 (45%)
α-Glucosidase inhibitors	40 (15%)

Table 1 continued

Variables	
Sulfonylureas	27 (10%)
Biguanides	131 (48%)
Glinides	27 (10%)
Thiazolidinediones	32 (12%)
Insulin	86 (31%)
GLP-1 receptor agonist	37 (13%)

Unless indicated otherwise, data are presented as mean ± standard deviation

HbA1c glycated hemoglobin, A1c, SGLT2 sodium glucose co-transporter 2, LDL low-density lipoprotein, HDL high-density lipoprotein, eGFR estimated glomerular filtration rate, DPP-4 dipeptidyl peptidase-4, GLP-1 glucagon-like peptide-1

Pharmacodynamic differences among SGLT2 inhibitors due to varying renal thresholds for glucose excretion and/or SGLT1/2 selectivity have been reported; thus, the glucose-lowering effects might also differ by type of SGLT2 inhibitor [6]. However, we did not observe any statistically significant differences in HbA1c levels after 3 months of treatment for the 6 different SGLT2 inhibitors in the ANOVA (Supplemental Fig. 1) nor in the multiple regression analyses adjusted for baseline eGFR (data not shown).

The glucose-lowering effect of SGLT2 inhibitors was greater when SGLT2 inhibitors were used in combination with biguanides when compared to using them without biguanides (without vs. with biguanides: $-0.31 \pm 0.91\%$ vs. $-0.55 \pm 0.94\%$; $p < 0.05$); however, there were no differences in the glucose-lowering effect of SGLT2 inhibitors between the groups with/without other antidiabetic drugs (Fig. 3). In addition, we found no differences in the glucose-lowering effect of SGLT2 inhibitors when SGLT2 inhibitors were used alone ($n = 37$) versus when they were used in combination with other antidiabetic drugs ($n = 238$) (monotherapy vs. combination therapy: $-0.33 \pm 0.97\%$ vs. $-0.44 \pm 0.93\%$; Fig. 3). The proportion of patients who achieved glycemic control of the HbA1c target of $< 7\%$ increased

Table 2 The effects of SGLT2 inhibitors on clinical parameters

Variables	Available data, <i>n</i>	Before the treatment	3 month after the treatment	<i>p</i> value
HbA1c (%)	275	7.8 ± 1.2	7.4 ± 1.0	< 0.0001
Rate of HbA1c < 7%, <i>n</i> (%)	275	76 (28%)	108 (39%)	< 0.0001
Random blood glucose (mg/dL)	271	166 ± 57	153 ± 50	0.0002
Body weight (kg)	242	82 ± 19	80 ± 18	< 0.0001
LDL cholesterol (mg/dL)	217	113 ± 33	110 ± 31	0.12
Triglyceride (mg/dL)	229	212 ± 297	203 ± 193	0.68
HDL cholesterol (mg/dL)	228	51 ± 15	52 ± 14	0.02
Hematocrit (%)	234	41 ± 5	43 ± 5	< 0.0001
Serum creatinine (mg/dL)	255	0.76 ± 0.39	0.79 ± 0.42	< 0.0001
eGFR (ml/min/1.73 m ²)	255	83 ± 26	80 ± 25	0.001
Systolic blood pressure (mmHg)	261	134 ± 16	131 ± 18	0.003
Diastolic blood pressure (mmHg)	261	80 ± 12	78 ± 12	0.019

Unless indicated otherwise, data are presented as mean ± standard deviation

HbA1c glycated hemoglobin, SGLT2 sodium glucose co-transporter 2, LDL low-density lipoprotein, HDL high-density lipoprotein cholesterol, eGFR estimated glomerular filtration rate




Fig. 2 Seasonal changes in HbA1c levels 3 months after the start of SGLT2 inhibitor treatment

from 28% to 39% within 3 months of SGLT2 inhibitor treatment (Table 2).

Effects of SGLT2 Inhibitors on the Lipid Profile and Renal Function

We noted a significant increase in HDL cholesterol levels (HDL cholesterol at 3 months—baseline HDL cholesterol: 1.2 ± 7.5 mg/dL, $p < 0.05$; Table 2) in response to treatment with SGLT2 inhibitors; however, no such effect was

observed for LDL cholesterol or triglyceride levels (Table 2).

We also detected significant increases in serum creatinine levels (serum creatinine at 3 months—baseline serum creatinine: 0.03 ± 0.16 mg/dL, $p < 0.0001$; Table 2) and decreases in eGFRs (eGFR at 3 months—baseline eGFR: -3.0 ± 29.8 mL/min/1.73 m², $p < 0.01$; Table 2); however, no cases of severe renal dysfunction were reported in this study.

Predictors of Decreased HbA1c Levels After SGLT2 Inhibitor Treatment

Single regression analysis showed that the magnitude of changes in HbA1c levels 3 months after the start of treatment (HbA1c at 3 months—baseline HbA1c) was positively associated with age ($p < 0.01$), indicating that younger patients showed greater decreases in HbA1c levels (Table 3). Changes in HbA1c levels 3 months after the start of treatment were also positively associated with changes in HbA1c levels 1 month after the start of treatment




Fig. 3 Glucose-lowering effect of SGLT2 inhibitors as monotherapy or in combination with other anti-diabetic drugs (* $p < 0.05$). Antidiabetic drugs included biguanides (BG), insulin, sulfonylureas (SU), DPP-4 inhibitors

(DPP-4i), glinides, α -glucosidase inhibitors (β -GI), GLP-1 receptor agonist (GLP-1RA), and thiazolidinediones (TZD)

(HbA1c at 1 months—baseline HbA1c) ($p < 0.001$). Moreover, we detected negative associations with baseline (pre-treatment) HbA1c levels ($p < 0.001$; Table 3; Fig. 4) and baseline eGFRs ($p < 0.001$; Table 3; Fig. 5). The greatest decrease in HbA1c levels associated with SGLT2 inhibitor treatment was observed among patients with an eGFR ≥ 100 mL/min/ 1.73 m 2 (Fig. 5). No association was found between changes in HbA1c levels and BMI (Table 3; Fig. 6).

We then performed a multiple regression analysis adjusting for age and sex to clarify the independent determinants for changes in HbA1c levels 3 months after the start of treatment (Table 3). This analysis showed that baseline HbA1c levels ($p < 0.0001$), baseline eGFR ($p < 0.05$), and changes in HbA1c levels 1 month after the start of treatment ($p < 0.001$)

were predictors for changes in HbA1c levels after 3 months of treatment. The multiple regression analysis adjusting for age, sex and concomitant antidiabetic drugs also showed that those three factors are predictors for changes in HbA1c levels after 3 months of treatment (Supplemental Table 1). These indicate that decreases in HbA1c levels were greater in patients with high baseline HbA1c levels, high baseline eGFRs, and greater initial responses to SGLT2 inhibitor treatment.




DISCUSSION

We show that SGLT2 inhibitors are effective in decreasing HbA1c levels after 3 months of treatment with no severe side effects. The predictors for the effects of SGLT2 inhibitor

Table 3 Determinants of change in HbA1c after 3 months treatment

Variables	Univariate		Multivariate			
	β (95% CI)	<i>p</i> value	Model 1		Model 2	
			β (95% CI)	<i>p</i> value	β (95% CI)	<i>p</i> value
Age	0.12 (0.03, 0.21)	< 0.01	0.04 (− 0.05, 0.12)	0.39	0.04 (− 0.07, 0.16)	0.48
Male	0.83 (− 0.46, 2.13)	0.20	0.59 (− 0.55, 1.73)	0.31	0.59 (− 0.74, 1.92)	0.38
BMI	− 0.04 (− 0.23, 0.16)	0.73				
Baseline HbA1c	− 0.43 (− 5.18, − 3.33)	< 0.001	− 4.14 (− 5.10, − 3.19)	< 0.0001		
Baseline eGFR	− 0.09 (− 0.14, − 0.04)	< 0.001			− 0.07 (− 0.13, − 0.01)	0.02
Change in HbA1c after 1 month	0.2 (0.05, 0.35)	< 0.001				0.89 (0.76, 1.02)
						< 0.001

BMI body mass index, HbA1c glycated hemoglobin, eGFR estimated glomerular filtration rate

**Fig. 4** Effects of SGLT2 inhibitor treatment on changes in HbA1c levels, stratified by baseline HbA1c levels**Fig. 5** Effects of SGLT2 inhibitor treatment on changes in HbA1c levels, stratified by estimated glomerular filtration rates (eGFR)**Fig. 6** Effects of SGLT2 inhibitor treatment on changes in HbA1c levels, stratified by baseline body mass index (BMI)

treatment were high baseline HbA1c levels, a high eGFR, and greater decreases in HbA1c levels after 1 month of treatment.

It has not yet been established which patients are ideal candidates for SGLT2 inhibitor treatment in terms of glucose lowering. The ASSIGN-K study showed that SGLT2 inhibitors were more effective in patients with high baseline HbA1c levels and a shorter duration of diabetes, while age, sex, and BMI were not

associated with a decrease in HbA1c levels 12 weeks after the start of ipragliflozin treatment [7]. Meta-regression analysis showed that high baseline glycemia increased the glucose-lowering effect of other oral agents such as sulfonylureas, glinides, biguanides, thiazolidinediones, α -glucosidase inhibitors, and DPP4 inhibitors [8–10]. Consistent with previous studies, our results show that decreased HbA1c levels are associated with baseline HbA1c levels. The fraction of filtered glucose excreted by SGLT2 inhibitors increases along with an increase in plasma glucose concentration [11]. This explains why the glucose-lowering effect of SGLT2 inhibitors was greater in subjects with high HbA1c levels when compared to those with low HbA1c levels.

It is known that SGLT2 inhibitors decrease body weight through caloric elimination, and that they are a suitable treatment option for obese diabetic patients [12]. However, our study shows that SGLT2 inhibitors decreased HbA1c levels regardless of baseline BMI, indicating that BMI is not a predictor of lowering HbA1c levels [13–14].

SGLT2 inhibitors stimulate food intake as a compensation for the energy loss due to glycosuria. It has been reported that SGLT2 inhibition may increase food intake in an animal model of obesity [15]. Thus, if patients quite caloric restriction, a rebound of hyperglycemia may occur. In addition, for antidiabetics such as sulfonylurea, rebound of the glucose-lowering effect or secondary failure due to desensitization of insulin secretion has been reported [16]. However, previous trials showed that SGLT2 inhibitors have a constant effect on lowering HbA1c levels without a glucose-lowering rebound [2, 17]. Consistent with earlier studies, we show that decreased HbA1c levels after 3 months of treatment were associated with decreased HbA1c levels after 1 month of treatment. This suggests that the 3-month glucose-lowering effect of SGLT2 inhibitor treatment can be predicted by the initial effects of the treatment.

It is known that SGLT2 inhibitors do not decrease HbA1c levels more than placebos in patients with moderate renal impairment of $\text{eGFR} < 45 \text{ mL/in}/1.73 \text{ m}^2$ [12]. However, it has

not been determined whether the glucose-lowering efficacy of SGLT2 can be predicted by baseline eGFR. We revealed that another predictor of SGLT2 inhibitor response is baseline high eGFR. SGLT2, which is located in the brush border membrane of the renal proximal tubule, is inhibited extracellularly from the lumen of the proximal tubule after SGLT2 inhibitors are filtered by the glomerulus [18]. Therefore, the glucose-lowering effect of SGLT2 inhibitors depends on the capacity of glucose filtration represented by the eGFR.

HbA1c values exhibited a seasonal change in Japanese patients with type 2 diabetes [19]. It has been reported that HbA1c values were highest in March and lowest in October, with a difference of 0.30% [20]. Consistent with this study, HbA1c levels were highest in February and March in our data. The reasons for the higher HbA1c levels during these months might be decreased physical activity due to cold weather and increased dietary calorie intake in winter [19].

When comparing the glucose-lowering effects of the 6 different SGLT inhibitors used in this study, no statistically significant difference was detected, although tofogliflozin had the greatest effect. The similarity in chemical structures of these SGLT2 inhibitors likely explains their class effects for glucose lowering [21, 22].

The glucose-lowering effect of SGLT2 inhibitors is independent of insulin, including beta-cell function and insulin sensitivity [23]. Therefore, the prevalence of hypoglycemia is rare, especially in the absence of insulin secretagogues and insulin. SGLT2 inhibitor-related serious adverse effects have been reported previously [24]. We experienced 10 cases of SGLT2 inhibitor-related adverse effects (itching, frequent urination, urinary tract infection, fainting, feeling thirsty, and lower limb numbness) leading to a discontinuation of SGLT2 inhibitor treatment (Fig. 1). SGLT2 inhibitor treatment was discontinued within 4 weeks in 9 of these cases. No cases of ketosis/ketoacidosis were observed, although urinary ketone tests were positive in 8 of 233 patients after SGLT2 inhibitor treatment (Supplemental Table 1). No study has thus far reported predictors for the adverse effects of SGLT2 inhibitors. Therefore, to avoid serious adverse effects, it is crucial to

identify suitable patients for SGLT2 inhibitor treatment by following the recommendation for the appropriate use of SGLT2 inhibitors by the Japanese Diabetes Society (<http://www.jds.or.jp>), and to carefully follow-up patients after administration of SGLT2 inhibitors, particularly during the first 4 weeks of treatment [24].

The present study has several limitations. First, we only analyzed patients who had been treated with SGLT2 inhibitors for at least 3 months at our hospitals; thus, our study was an inherent patient selection bias. Second, this was a retrospective study with a small sample size. Third, the observation period was relatively short; therefore, we could not analyze the long-term effects of SGLT2 inhibitor treatment. Fourth, 87% of the patients were undergoing combination therapy with SGLT2 inhibitors and other antidiabetic drugs. Thus, we could not exclude the effects of concomitant antidiabetic drugs. Large clinical cohort studies with a longer observation period are needed to clarify the reliable predictors in type 2 diabetes patients of the effect of SGLT2 inhibitors for lowering HbA1c levels.

CONCLUSIONS

In conclusion, we show that high baseline HbA1c levels and a high baseline eGFR were predictors for the effect of SGLT2 inhibitors of lowering HbA1c levels after 3 months of treatment. Moreover, the effect of SGLT2 inhibitors could be predicted by the level of decrease in HbA1c levels after 1 month of treatment (i.e., the initial response to treatment). This suggests that patients with type 2 diabetes who receive the greatest glucose-lowering effect with SGLT2 inhibitor treatment are those with preserved renal function and high HbA1c levels regardless of BMI. Moreover, SGLT2 inhibitor treatment efficacy can be predicted by the patients' initial response to treatment.

ACKNOWLEDGEMENTS

This work was partially supported by JSPS Kakenhi Grants (Grant numbers 16H05299 and

26248050), GSK Japan Research Grant, The Uehara Memorial Foundation, and Takeda Science Foundation. Article processing charges were funded by the authors. We thank the staff of the Hospital Information Center at Tokushima University Hospital and Masato Mori for extracting clinical data from medical records. We would like to thank Editage (<http://www.editage.jp>) for English language editing. All named authors meet the International Committee of Medical Journal Editors (ICMJE) criteria for authorship for this manuscript, take responsibility for the integrity of the work as a whole, and have given final approval for the version to be published.

Disclosures. M. Matsuhisa received research funding and lecture fees from Sanofi, Takeda Pharmaceutical Company Ltd., Eli-Lilly, Novo Nordisk Pharma, Astellas Pharma Inc., Boehringer Ingelheim, Daiichi Sankyo Co. Ltd., Tanabe-Mitsubishi Pharma, Novartis Pharmaceuticals Corporation. M. Shimabukuro received research funding from AstraZeneca and lecture fees from Astellas Pharma Inc., Ono Pharmaceutical Company Ltd, Taisho Toyama Pharmaceutical Co., Ltd., Kowa Company, Ltd., Nippon Boehringer Ingelheim, AstraZeneca, and Daiichi Sankyo Co. Ltd. M. Sata received research funding from Tanabe-Mitsubishi, Takeda, Astellas, Bayer Healthcare, Daiichi-Sankyo, MSD, and Ono, and lecture fees from Astellas, Boehringer Ingelheim, Bayer Healthcare, Mochida, Takeda, Tanabe-Mitsubishi, Novartis, AstraZeneca, MSD, and Shionogi. The Department of Cardio-Diabetes Medicine, Tokushima University Graduate School, is supported in part by unrestricted research grants from Actelion, Boehringer Ingelheim, Kowa, and Tanabe-Mitsubishi. Shusuke Yagi, Ken-ichi Aihara, Takeshi Kondo, Kiyoe Kurahashi, Sumiko Yoshida, Itsuro Endo, Daiju Fukuda, Yutaka Nakaya, Kin-ichiro Suwaki, Takashi Takeji, Toshihiro Wada, Hotimah M. An Salim, Saori Hama, Tomomi Matsura, Takayuki Ise, Kenya Kusunose, Koji Yamaguchi, Takeshi Tobiume, Hirotugu Yamada, Takeshi Soeki, Tetsuzo Wakatsuki and Masashi Akaiki have nothing to disclose.

Compliance with Ethics Guidelines. All procedures were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1964, as revised in 2013. The study protocol was approved by the Ethics Committee of Tokushima University Hospital (No. 2432) and Shikoku Central Hospital (No. 25).

Data Availability. The datasets during and/or analyzed during the current study are available from the corresponding author on reasonable request.

REFERENCES

1. Tanaka A, Node K. Emerging roles of sodium-glucose cotransporter 2 inhibitors in cardiology. *J Cardiol.* 2017;69:501–7.
2. Zinman B, Wanner C, Lachin JM, et al. Empagliflozin, cardiovascular outcomes, and mortality in type 2 diabetes. *N Engl J Med.* 2015;373:2117–28.
3. Inzucchi SE, Zinman B, Wanner C, et al. SGLT-2 inhibitors and cardiovascular risk: proposed pathways and review of ongoing outcome trials. *Diabetes Vasc Dis Res.* 2015;12:90–100.
4. Salim HM, Fukuda D, Yagi S, Soeki T, Shimabukuro M, Sata M. Glycemic control with ipragliflozin, a novel selective SGLT2 inhibitor, ameliorated endothelial dysfunction in streptozotocin-induced diabetic mouse. *Front Cardiovasc Med.* 2016;3:43.
5. American Diabetes A. Standards of medical care in diabetes—2016 abridged for primary care providers. *Clin Diabetes.* 2016;34:3–21.
6. Sha S, Polidori D, Farrell K, et al. Pharmacodynamic differences between canagliflozin and dapagliflozin: results of a randomized, double-blind, crossover study. *Diabetes Obes Metab.* 2015;17:188–97.
7. Iemitsu K, Iizuka T, Takihata M, et al. Factors influencing changes in hemoglobin A1c and body weight during treatment of type 2 diabetes with ipragliflozin: interim analysis of the ASSIGN-K study. *J Clin Med Res.* 2016;8:373–8.
8. Kalra S. Sodium glucose co-transporter-2 (SGLT2) inhibitors: a review of their basic and clinical pharmacology. *Diabetes Ther Res Treat Educ Diabetes Relat Disord.* 2014;5:355–66.
9. Bloomgarden ZT, Dodis R, Viscoli CM, Holmboe ES, Inzucchi SE. Lower baseline glycemia reduces apparent oral agent glucose-lowering efficacy: a meta-regression analysis. *Diabetes Care.* 2006;29:2137–9.
10. Yagi S, Aihara K, Akaike M, et al. Predictive factors for efficacy of dipeptidyl peptidase-4 inhibitors in patients with type 2 diabetes mellitus. *Diabetes Metab J.* 2015;39:342–7.
11. Abdul-Ghani MA, DeFronzo RA, Norton L. Novel hypothesis to explain why SGLT2 inhibitors inhibit only 30–50% of filtered glucose load in humans. *Diabetes.* 2013;62:3324–8.
12. Kohan DE, Fioretto P, Tang WH, List JF. Long-term study of patients with type 2 diabetes and moderate renal impairment shows that dapagliflozin reduces weight and blood pressure but does not improve glycemic control. *Kidney Int.* 2014;85:962–71.
13. Rosenstock J, Jelaska A, Frappin G, et al. Improved glucose control with weight loss, lower insulin doses, and no increased hypoglycemia with empagliflozin added to titrated multiple daily injections of insulin in obese inadequately controlled type 2 diabetes. *Diabetes Care.* 2014;37:1815–23.
14. Sakai S, Kaku K, Seino Y, et al. Efficacy and safety of the SGLT2 inhibitor luseogliflozin in Japanese patients with type 2 diabetes mellitus stratified according to baseline body mass index: pooled analysis of data from 52-Week phase III trials. *Clin Ther.* 2016;38(843–62):e9.
15. Han S, Hagan DL, Taylor JR, et al. Dapagliflozin, a selective SGLT2 inhibitor, improves glucose homeostasis in normal and diabetic rats. *Diabetes.* 2008;57:1723–9.
16. Satoh J, Takahashi K, Takizawa Y, et al. Secondary sulfonylurea failure: comparison of period until insulin treatment between diabetic patients treated with gliclazide and glibenclamide. *Diabetes Res Clin Pract.* 2005;70:291–7.
17. Maegawa H, Tobe K, Tabuchi H, Nakamura I. Baseline characteristics and interim (3-month) efficacy and safety data from STELLA-LONG TERM, a long-term post-marketing surveillance study of ipragliflozin in Japanese patients with type 2 diabetes in real-world clinical practice. *Expert Opin Pharmacother.* 2016;17:1985–94.
18. Ghezzi C, Hirayama BA, Gorraitz E, Loo DD, Liang Y, Wright EM. SGLT2 inhibitors act from the extracellular surface of the cell membrane. *Physiol Rep.* 2014;2:e12058.
19. Ishii H, Suzuki H, Baba T, Nakamura K, Watanabe T. Seasonal variation of glycemic control in type 2 diabetic patients. *Diabetes Care.* 2001;24:1503.

20. Furuta M, Tomisaka R, Yamana A, et al. Evaluation of seasonal changes in hemoglobin A1c in diabetic patients. *Rinsho byori Jpn J Clin Pathol*. 2012;60:599–604.
21. Yanai H, Katsuyama H, Hamasaki H, et al. Sodium-glucose cotransporter 2 inhibitors: possible anti-atherosclerotic effects beyond glucose lowering. *J Clin Med Res*. 2016;8:10–4.
22. Mosley JF 2nd, Smith L, Everton E, Fellner C. Sodium-glucose linked transporter 2 (SGLT2) inhibitors in the management of type-2 diabetes: a drug class overview. *P T*. 2015;40:451–62.
23. Clar C, Gill JA, Court R, Waugh N. Systematic review of SGLT2 receptor inhibitors in dual or triple therapy in type 2 diabetes. *BMJ Open*. 2012;2:e001007.
24. Yabe D, Nishikino R, Kaneko M, Iwasaki M, Seino Y. Short-term impacts of sodium/glucose co-transporter 2 inhibitors in Japanese clinical practice: considerations for their appropriate use to avoid serious adverse events. *Expert Opin Drug Saf*. 2015;14:795–800.

paper 4

ORIGINALITY REPORT

24%

SIMILARITY INDEX

21%

INTERNET SOURCES

26%

PUBLICATIONS

12%

STUDENT PAPERS

PRIMARY SOURCES

1

e-sciencecentral.org

Internet Source

6%

2

dmsjournal.biomedcentral.com

Internet Source

3%

3

Shusuke Yagi, Daiju Fukuda, Ken-ichi Aihara, Masashi Akaike, Michio Shimabukuro, Masataka Sata. "n-3 Polyunsaturated Fatty Acids: Promising Nutrients for Preventing Cardiovascular Disease", Journal of Atherosclerosis and Thrombosis, 2017

Publication

1%

4

doaj.org

Internet Source

1%

5

Alessandro Ciresi, Enrica Vigneri, Stefano Radellini, Felicia Pantò, Carla Giordano. "Liraglutide Improves Cardiovascular Risk as an Add-on to Metformin and Not to Insulin Secretagogues in Type 2 Diabetic Patients: A Real-life 48-Month Retrospective Study", Diabetes Therapy, 2017

1%

6

eprints.soton.ac.uk

Internet Source

1%

7

Paul Chi Ho Lee, Yunjuan Gu, Man Yi Yeung, Carol Ho Yi Fong, Yu Cho Woo, Wing Sun Chow, Kathryn Tan, Karen Siu Ling Lam.

"Dapagliflozin and Empagliflozin Ameliorate Hepatic Dysfunction Among Chinese Subjects with Diabetes in Part Through Glycemic Improvement: A Single-Center, Retrospective, Observational Study", Diabetes Therapy, 2018

Publication

1%

8

Yagi, Shusuke, Ken-ichi Aihara, Masashi Akaike, Daiju Fukuda, Hotimah Masdan Salim, Masayoshi Ishida, Tomomi Matsuura, Takayuki Ise, Koji Yamaguchi, Takashi Iwase, Hirotsugu Yamada, Takeshi Soeki, Tetsuzo Wakatsuki, Michio Shimabukuro, Toshio Matsumoto, and Masataka Sata. "Predictive Factors for Efficacy of Dipeptidyl Peptidase-4 Inhibitors in Patients with Type 2 Diabetes Mellitus", Diabetes & Metabolism Journal, 2015.

Publication

1%

9

Shusuke Yagi, Daisuke Kondo, Takayuki Ise, Daiju Fukuda et al. "Association of Decreased Docosahexaenoic Acid Level After Statin Therapy and Low Eicosapentaenoic Acid Level

1%

with In-Stent Restenosis in Patients with Acute Coronary Syndrome", Journal of Atherosclerosis and Thrombosis, 2019

Publication

10

Noriko Sugasawa, Ayako Katagi, Hirotsugu Kurobe, Taisuke Nakayama et al. "Inhibition of Atherosclerotic Plaque Development by Oral Administration of α -Glucosyl Hesperidin and Water-Dispersible Hesperetin in Apolipoprotein E Knockout Mice", Journal of the American College of Nutrition, 2018

Publication

1%

11

kundoc.com

Internet Source

1%

12

Gopalakrishnan, Sathej, Amit Khatri, Sven Mensing, Rebecca Redman, Rajeev Menon, and Jiahong Zha. "Exposure–Response Relationship for Ombitasvir and Paritaprevir/Ritonavir in Hepatitis C Virus Subgenotype 1b-Infected Japanese Patients in the Phase 3 Randomized GIFT-I Study", Advances in Therapy, 2016.

Publication

1%

13

www.biomedcentral.com

Internet Source

1%

14

Hiromu Yamazaki, Shusuke Yagi, Yuta Torii, Rie Amano et al. "Edoxaban improves acute

1%

venous thromboembolism while preserving protein C and protein S levels", Journal of Cardiology, 2018

Publication

15

"Poster Presentation", Journal of Diabetes Investigation, 2012.

Publication

1%

16

www.ncbi.nlm.nih.gov

Internet Source

1%

17

Antonella Napolitano, Sam Miller, Peter R. Murgatroyd, Elizabeth Hussey et al. "Exploring glycosuria as a mechanism for weight and fat mass reduction. A pilot study with remogliflozin etabonate and sergliflozin etabonate in healthy obese subjects", Journal of Clinical & Translational Endocrinology, 2014

Publication

1%

18

Eugene Han, Ari Kim, Sung Jae Lee, Je-Yon Kim, Jae Hyeon Kim, Woo Je Lee, Byung-Wan Lee. "Characteristics of Dapagliflozin Responders: A Longitudinal, Prospective, Nationwide Dapagliflozin Surveillance Study in Korea", Diabetes Therapy, 2018

Publication

1%

19

Abdul-Ghani, M. A., R. A. DeFronzo, and L. Norton. "Novel Hypothesis to Explain Why SGLT2 Inhibitors Inhibit Only 30-50% of

1%

Filtered Glucose Load in Humans", Diabetes, 2013.

Publication

20

Ivano Riva, Andreas Katsanos, Francesco Oddone, Luciano Quaranta. "The Effect of Phacoemulsification on Intraocular Pressure in Eyes with Hyperfiltration Following Trabeculectomy: A Prospective Study", Advances in Therapy, 2017

1%

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%